

AGRI-PLEX[®] CROP KICKER

SUPPLEMENTAL PLANT FOOD FOR FOLIAR APPLICATION

12-4-6

-GUARANTEED ANALYSIS-

Total Nitrogen (N)	12.00%
1.20% Ammoniacal Nitrogen (N)	
1.78% Nitrate Nitrogen (N)	
9.02% Urea Nitrogen (N)	
Available Phosphate (P ₂ O ₃)	4.00%
Soluble Potash (K ₂ O)	6.00%
Boron (B)	0.02%
Copper (Cu)	0.05%
Iron (Fe)	0.10%
0.05% Chelated Iron (Fe)	
Manganese (Mn)	0.05%
0.05% Chelated Manganese (Mn)	
Zinc (Zn)	0.05%
0.05% Chelated Zinc (Zn)	

Derived from: Ammonium 7um Polyphosphate, Urea, Potassium, Nitrate, Zinc EDTA, Manganese EDTA, Iron EDTA and Copper EDTA and Sodium Borate.

DIRECTIONS FOR USE

Crop Kicker will disperse in water with minimum agitation. It is compatible with most pesticides. Always check compatibility before mixing. Follow the mixing sequence: 1. Water, 2. Crop Kicker, 3. Pesticide –Minimum GPA: 20 ground, 5 air.. For best results apply early morning or late afternoon when leaf pores are open. Spray as fine mist during periods of little dew to avoid runoff of leaf surfaces.

INFORMATION

Crop Kicker is a liquid plant food compound for foliar application containing the primary elements and micronutrients. It is recommended for use as a supplement to a conventional fertility program—potentially enhancing yield and/or quality. Crop Kicker supplements soil fertilization by supplying essential nutrients when plant food need are highest—at flowering and/or fruiting stages or in stress conditions. Crop Kicker is specially formulated for cotton, peanuts, soybeans and vegetables but can be used for most crops.

RECOMMENDED RATES

Cotton: Apply one quart per acre early season. Increase to 2-4 quarts per acre for mid or late season or by first bloom and repeat at 10-14 day intervals.
Peanuts: Use one quart per acre early season. Increase rate up to 2-4 quarts per acre by first bloom and continue at 10-14 day intervals.
Soybean: Use 2-4 quarts per acre with each application starting no later than first bloom. Continue as needed through pod development.
Vegetables: Use 1-4 quarts per acre starting depending on size of plants. First application should be made prior to flowering and/or fruiting. Continue throughout season at 7-10 day intervals.
Other Crops: Use 1-4 quarts per acre starting no later than first bloom or fruiting stage and continue every 10-14 days along with regular spray program.

CONDITIONS OF SALE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. Since weather, crop, soil and other conditions may vary, Agri-Business Technologies, Inc. and/or the seller make no warranty of any kind, expressed or implied, concerning the use of this product. The user assumes all risks of use or handling whether or not in accordance with directions or suggestions.

STORAGE

Store in a cool place in original container in a locked storage area inaccessible to children, pets, or wildlife.

CONTAINER DISPOSAL

Triple rinse empty containers and offer for recycling or dispose in accordance with federal, state and local authorities.

FIRST AID

If Swallowed: DO NOT INDUCE VOMITING. Immediately call physician
Eye Contact: Flush eyes with clean water for 15 minutes. Immediately call a physician.
Skin Contact: Remove contaminated clothing and wash skin with soap and water. Call a physician if irritation develops or persists.
If Inhaled: Move to fresh air and call a physician if irritation develops or persists.

If there is a contact made with the spray solution containing pesticide and/or adjuvants, follow the "Statement of Practical Treatment" on the product labeling.

CAUTION

Keep Out Of Reach Of Children

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin and clothing. Avoid breathing spray mist. READ LABEL! FOLLOW DIRECTIONS!

Manufactured For:

Agri-Business Technologies, Inc.

1102 3rd Avenue • Albany GA 31707

Net Weight: 50 lbs. – 22.68 kg

F1285